

daflon[®] 500 mg

達夫隆

Micronized purified flavonoid fraction

Presentation and composition

Box of 30 film-coated tablets.

Micronized purified flavonoid fraction corresponding to:

- diosmin: 90 %
- flavonoids expressed as hesperidin: 10 %

Excipients q.s. for one tablet.

Therapeutic indications

This medicine is a venotonic (it increases venous tone) and a vasculoprotector (it increases resistance in small blood vessels).

It is recommended for treating venous circulation disorders (swollen legs, pain, restless legs) and for treating symptoms due to acute hemorrhoidal attack.

Side effects

Some cases of routine gastric disorders and neuro-vegetative disorders (feeling of discomfort) have been reported, not requiring discontinuation of treatment.

Report any untoward or undesirable effect which has not been mentioned in this package insert to your doctor or your pharmacist.

Precautions

Acute hemorrhoidal attack:

If the hemorrhoid symptoms do not disappear within 15 days, you should ask your doctor for advice.

Venous circulation disorders:

The most effective way of giving this treatment is in combination with a healthy lifestyle. Avoid exposure to the sun, heat, excessive standing and being overweight. Walking and wearing special support stockings stimulate blood circulation.

If you are in any doubt, do not hesitate to ask your doctor or your pharmacist for advice.

Interactions with other medicines and other forms of interactions

You must always tell your doctor or your pharmacist about any other treatment you are receiving.

Pregnancy – Breast-Feeding

Inform your doctor if you become pregnant.

In the absence of data on entering breast milk, breast-feeding should be avoided during treatment.

In general, if you are pregnant or breast-feeding, you should always ask your doctor or pharmacist for advice before using a medication.

Dosage and method of administration

Oral route.

The tablets should be taken at meal times.

Venous insufficiency:

2 tablets daily, one at midday and one in the evening.

Acute hemorrhoidal attack:

A 4-day course of 6 tablets daily, followed by 4 tablets daily over the next 3 days.

In all cases strictly comply with your doctor's prescription.

Storage conditions

Do not use after the expiry date indicated on the outer packaging.



LES LABORATOIRES SERVIER – FRANCE

Manufacturer:

LES LABORATOIRES SERVIER

INDUSTRIE

45520 Gidy – France

daflon[®] 500 mg

達夫隆

Micronized purified flavonoid fraction

包裝和成份：

每盒 30 片包衣藥片。每片藥片含微粒化，純化的黃酮類成份如下：

- Diosmin 90%
- Hesperidin 10%

適應症：

Daflon[®] 500mg 能加強靜脈張力和具有血管保護作用。Daflon[®] 500mg 推薦用于治療靜脈循環系統障礙（症狀如：下肢腫脹，疼痛，沉重感及腿部不適等等），亦用于治療急性痔瘡發作。

副作用：

少數病人出現輕微的胃腸道不適及植物神經功能紊亂，但并不需要停止治療。

如有其他副作用出現，請通知您的醫生或藥劑師。

注意事項：

性痔瘡發作：

如症狀持續 15 天仍未消失，請通知您的醫生。

靜脈循環系統障礙：

本藥治療期間如養成平衡良好的生活方式將獲得最大效益。

如避免暴露于陽光，高熱的地方。避免長時間的站立及避免體重超重。適量地步行及穿著彈力襪以改善血液循環。

藥物相互作用：

請通知您的醫生或藥劑師您正在服用的其他藥物。

妊娠和授乳：

請通知您的醫生如果您將懷孕。

由于缺少授乳的資料，正在接受治療的期間應避免授乳。

一般而言，如懷孕或授乳都應先徵詢醫生的意見才服用任何藥物。

服法及用量：

口服，于用餐時服用。

治療靜脈功能不全：

每天 2 片，午餐及晚餐時各服一片。

急性痔瘡發作：

首 4 日，每天 6 片然後每天 4 片服用 3 天。

請遵照醫生指示服用。

貯存條件：

請依照盒上限期前服用。



Les Laboratoires Servier – France

Manufacturer:

Les Laboratoires Servier Industrie

45520 GIDY – FRANCE